


Contents lists available at [ScienceDirect](http://ScienceDirect.com)

Taiwanese Journal of Obstetrics & Gynecology

journal homepage: www.tjog-online.com

Case Report

Prenatal diagnosis and molecular cytogenetic characterization of a *de novo* unbalanced reciprocal translocation of der(9)t(9;14)(p24.2;q32.11) associated with 9p terminal deletion and 14q distal duplicationChih-Ping Chen^{a, b, c, d, e, f, *}, Chen-Ju Lin^{a, g}, Schu-Rern Chern^b, Peih-Shan Wu^h, Yen-Ni Chen^a, Shin-Wen Chen^a, Chen-Chi Lee^a, Li-Feng Chen^a, Chien-Wen Yang^b, Wayseen Wang^{b, i}^a Department of Obstetrics and Gynecology, MacKay Memorial Hospital, Taipei, Taiwan^b Department of Medical Research, MacKay Memorial Hospital, Taipei, Taiwan^c Department of Biotechnology, Asia University, Taichung, Taiwan^d School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan^e Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan^f Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan^g Department of Medicine, MacKay Medical College, New Taipei City, Taiwan^h Gene Biodesign Co. Ltd, Taipei, Taiwanⁱ Department of Bioengineering, Tatung University, Taipei, Taiwan

ARTICLE INFO

Article history:

Accepted 25 May 2016

Keywords:

9p deletion syndrome
14q duplication syndrome
partial monosomy 9p
partial trisomy 14q
prenatal diagnosis

ABSTRACT

Objective: We present molecular cytogenetic characterization of a prenatally detected derivative chromosome 9 [der(9)] of unknown origin.**Case Report:** A 35-year-old woman underwent amniocentesis at 18 weeks of gestation because of advanced maternal age, which revealed a der(9) chromosome of unknown origin. The parental karyotypes were normal. Array comparative genomic hybridization (aCGH) analysis revealed a 2.593 Mb deletion of 9p24.3–p24.2 encompassing *DOCK8*, *KANK1*, *DMRT1*, and *VLDLR* and a 16.65 Mb duplication of 14q32.11–q32.33 encompassing *DLK1*, *RTL1*, *MEG3*, *RTL1as*, and *MEG8*. Quantitative fluorescent polymerase chain reaction (QF-PCR) analysis using D9S937 (9p24.2) and D14S605 (14q32.2) showed a paternal origin of 9p24.2 deletion and a paternal origin of 14q32.2 duplication consistent with a paternal origin of the *de novo* aberrant chromosome of der(9)t(9p;14q). The fetal karyotype was 46,XX,der(9)t(9;14)(p24.2;q32.11). Metaphase fluorescence *in situ* hybridization (FISH) analysis using RP11-57K23 (14q32.33), RP11-31F19 (9p24.3), RP11-30O14 (9p21.1), and RP11-1105I14 (14q11.2) confirmed an unbalanced reciprocal translocation of der(9)t(9p;14q). We discuss 9p deletion syndrome and 14q duplication syndrome in this case.**Conclusion:** Molecular cytogenetic techniques such as aCGH, FISH, and QF-PCR are useful in the determination of the origin and nature of a prenatally detected *de novo* derivative chromosome of unknown origin.Copyright © 2016, Taiwan Association of Obstetrics & Gynecology. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Prenatal diagnosis of *de novo* chromosome aberration of unknown origin may require molecular cytogenetic characterizationsuch as array comparative genomic hybridization (aCGH), fluorescence *in situ* hybridization (FISH), and quantitative fluorescent polymerase chain reaction (QF-PCR) to delineate the nature and the parental origin of the aberration [1].Here, we present our experience of molecular cytogenetic characterization of a prenatally detected derivative chromosome 9 [der(9)] of unknown origin by the use of aCGH, FISH, and QF-PCR, and the result was a *de novo* unbalanced reciprocal translocation* Corresponding author. Department of Obstetrics and Gynecology, MacKay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei 10449, Taiwan.
E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).

of der(9)t(9;14) (p24.2;q32.11) of paternal origin. Our presentation demonstrates the usefulness of molecular cytogenetic techniques such as aCGH, FISH, and QF-PCR in the determination of the origin and nature of a prenatally detected *de novo* derivative chromosome of unknown origin.

Case Report

A 35-year-old, gravid 2, para 1, woman underwent amniocentesis at 18 weeks of gestation because of advanced maternal age. Her husband was 40 years of age. The couple had a healthy 4-year-old boy, and there was no family history of congenital malformations. Amniocentesis revealed a der(9) with a deletion of terminal 9p and an addendum of an unknown segment at the breakpoint of 9p. The parental karyotypes were normal. aCGH analysis on cultured amniocytes by CytoScan 750K Array (Affymetrix, Santa Clara, CA, USA) showed the result of arr 9p24.3p24.2 (208,454–2,801,321) \times 1.0, 14q32.11q32.33 (90,637,774–107,284,437) \times 3.0, indicating a 2.593 Mb deletion of 9p24.3–p24.2 encompassing 16 genes including 8 Online Mendelian Inheritance in Man (OMIM) genes of *DOCK8*, *KANK1*, *DMRT1*, *DMRT3*, *DMRT2*, *SMARCA2*, *VLDLR*, and *KCNV2*, and a 16.65 Mb duplication of 14q32.11–q32.33 encompassing 519 genes including 142 OMIM genes that contain the paternally expressed genes (PEGs) of *DLK1* and *RTL1*, and the maternally expressed genes (MEGs) of *MEG3*, *RTL1as*, and *MEG8* (Figure 1). Metaphase FISH analysis on cultured amniocytes using the bacterial artificial chromosome probes of RP11-57K23 (14q32.33, 106,882,918–106,976,919, dye: FITC, spectrum green) [hg 19], RP11-31F19 (9p24.3, 547,217–692,143, dye: Texas Red, spectrum red) [hg 19], RP11-30014 (9p21.1, 28,765,550–28,966,389, dye: Cy5, spectrum yellow) [hg 19], and RP11-1105114 (14q11.2, 19,662,704–19,861,854, dye: DEAC, spectrum blue) [hg 19] showed the presence of one blue signal and one green signal in the normal chromosome 14, one yellow signal and one red signal in the normal chromosome 9, and one yellow signal and one green signal in the der(9) chromosome, indicating an unbalanced reciprocal translocation of der(9)t(9p;14q) (Figure 2). The fetal karyotype was 46,XX,der(9)t(9;14)(p24.2;q32.11) (Figure 3). Prenatal ultrasound findings were unremarkable. The pregnancy was subsequently terminated at 22 weeks of gestation, and a 408 g fetus was delivered with mild facial dysmorphism of micrognathia and low-set ears. The female external genitalia were normal. QF-PCR analysis on the DNAs extracted from the placental tissue and the parental bloods using informative markers of D9S937 (9p24.2), D9S2149 (9p21.1), D14S140 (14q31.3), and D14S605 (14q32.2) showed only one maternal allele at D9S937 (9p24.2) and a paternal allele:maternal allele ratio of 2:1 at D14S605 (14q32.2) in the fetus, indicating a paternal origin of the der(9) chromosome (Figure 4).

Discussion

The present case had a 2.593 Mb deletion of 9p24.2→pter encompassing the critical genes of *DOCK8*, *KANK1*, *DMRT1*, and *VLDLR* that may be responsible for the chromosome distal 9p deletion syndrome (OMIM 158170). Chromosome distal 9p deletion syndrome is characterized by major clinical features of mental retardation, hypotonia, seizures, trigonocephaly, synophrys, mid-face hypoplasia, short nose, depressed nasal bridge, anteverted nares, hypertelorism, up-slanting palpebral fissures, long philtrum, microstomia, high and narrow palate, small posteriorly rotated ears, hypoplastic labia majora, prominent labia minora and clitoris, cryptorchidism, hypospadias, wide-set nipples, hernias, scoliosis, diastasis recti, short and broad distal phalanges of fingers, square-shaped nails, and foot position anomalies; and relatively rare

malformations of congenital heart defects, microphthalmia, choanal atresia, stenotic external ear canals, cleft palate, diaphragmatic hernia, hydronephrosis, hypoplasia of the corpus callosum, enlarged cisterna magna, and postaxial hexadactyly of the fingers [2].

A 9p24.3 deletion is associated with 46,XY sex reversal (SRXY4; OMIM 154230) because the *DMRT* cluster of *DMRT1*, *DMRT2*, and *DMRT3* is located at 9p24.3, and *DMRT1* is the strongest candidate gene for the sex reversal [3–6]. *DMRT1* (OMIM 602424) encodes a male-specific transcriptional regulator of doublesex- and MAB3-related transcription factor 1 (*DMRT1*) that is involved in sex determination and differentiation [7,8]. *DMRT1* suppresses female differentiation in testes, and involves inhibition of meiosis in testes [3–5]. Haploinsufficiency of *DMRT1* will cause 46,XY gonadal dysgenesis and 46,XY ovotesticular disorder of sexual development [6]. Heterozygous disruption of *DOCK8* (OMIM 611432) either by a deletion or by a translocation breakpoint is associated with autosomal dominant mental retardation 2 (MRD2; OMIM 614113) [9]. Deletion of *KANK1* (OMIM 607704) causes familial cerebral palsy (cerebral palsy spastic quadriplegic 2; OMIM 612900) which is associated with parent-of-origin-dependent inheritance, and the affected individuals inherit the deletion of *KANK1* from paternal origin [10]. *VLDLR* encodes very low-density lipoprotein receptor (*VLDLR*) and is involved in Reelin signaling pathway and neuronal migration. Mutations of *VLDLR* (OMIM 192977) cause abnormal gyration, psychiatric disorders, and autosomal recessive cerebellar ataxia, mental retardation and dysequilibrium syndrome 1 (CAMRQ1; OMIM 224050) [11–14].

To date, at least 10 cases with fetal distal 9p deletion detected by prenatal invasive diagnostic procedures have been reported because of various indications such as abnormal maternal serum biochemistry, increased nuchal translucency, ambiguous external genitalia, male-to-female sex reversal, intrauterine growth restriction, advanced maternal age, familial chromosomal translocation and fetal structural abnormalities [2,15–22]. Stumm et al [15] reported prenatal diagnosis of mosaic r(9)(p24q34) in a male fetus because of the fetal ultrasound finding of male-to-female sex reversal. Vialard et al [16] reported prenatal diagnosis of del(9)(p22) in a male fetus with ambiguous external genitalia and in a female fetus associated with hypoplastic left heart and a single umbilical artery on fetal ultrasound. Witters et al [17] reported prenatal diagnosis of der(9)t(3;9)(p14.2;p24) in a male fetus associated with sex reversal and multiple anomalies on fetal ultrasound. Brisset et al [18] reported prenatal diagnosis of der(9)t(9;17)(p24.3;q24.3) in a female fetus associated with increased nuchal translucency, a single umbilical artery and partial agenesis of the cerebellar vermis on fetal ultrasound. Chen et al [19] reported prenatal diagnosis of mosaic r(9)(p24q34.3) in a male fetus with abnormal maternal serum screening and ambiguous external genitalia on fetal ultrasound. Chen et al [20] reported prenatal diagnosis of inv dup del(9) (:p22.1→p24.3::p24.3→qter) in a male fetus with ventriculomegaly on fetal ultrasound. Chen et al [2] reported prenatal diagnosis of 46,XX,del(9)(p24.1p24.3) because of abnormal maternal serum screening. Hou et al [21] reported prenatal diagnosis of del(9)(p13) in a female fetus associated with symbrachydactyly and omphalocele on fetal ultrasound. Penacho et al [22] reported prenatal diagnosis of r(9)(p24q34) in a female fetus associated with intrauterine growth restriction and slight craniofacial dysmorphism. The peculiar aspect of the present case is the diagnostic indication of advanced maternal age for amniocentesis with a result of a *de novo* unbalanced reciprocal translocation of der(9)t(9;14)(p24.2;q32.11) of paternal origin. This information is very important in the genetic counseling for the pathogenesis of a *de novo* chromosome aberration.

The 14q duplication syndrome involving dup 14q (14q22→q32) is characterized by high forehead, wide sutures and fontanelles,




Figure 1. aCGH on cultured amniocytes shows a 2.593 Mb deletion of 9p24.3–p24.2 encompassing *DOCK8*, *KANK1*, *DMRT1*, and *VLDLR*, and a 16.65 Mb duplication of 14q32.11–q32.33 encompassing *DLK1*, *RTL1*, *MEG3*, *RTL1as*, and *MEG8*. (A, C) chromosome 9 zoom-in view; (B, D) chromosome 14 zoom-in view. aCGH = array comparative genomic hybridization.




Figure 2. Metaphase FISH analysis on cultured amniocytes using the bacterial artificial chromosome probes of RP11-57K23 (14q32.33, 106,882,918–106,976,919, dye: FITC, spectrum green) [hg 19], RP11-31F19 (9p24.3, 547,217–692,143, dye: Texas Red, spectrum red) [hg 19], RP11-30014 (9p21.1, 28,765,550–28,966,389, dye: Cy5, spectrum yellow) [hg 19], and RP11-1105114 (14q11.2, 19,662,704–19,861,854, dye: DEAC, spectrum blue) [hg 19] shows the presence of one blue signal and one green signal in the normal chromosome 14, one yellow signal and one red signal in the normal chromosome 9, and one yellow signal and one green signal in the der(9) chromosome, indicating an unbalanced reciprocal translocation of der(9)t(9p;14q). FISH = fluorescence in situ hybridization.

sparse hair and eyebrows, board nasal bridge, hypertelorism, thin upper lip, abnormal ears, broad mouth, micrognathia, developmental delay, mild mental retardation, spasticity, hyperreflexia, premature puberty, and primordial short stature [23–27]. The 14q

duplication syndrome involving dup 14q (14q31→qter) is characterized by an asymmetric face, frontal bossing, low-set and dysplastic ears, hypertelorism, sparse eyebrows and lashes, prominent nasal bridge, small mandible, growth retardation, and moderate mental retardation [28–36].

The present case had a 16.65 Mb duplication of 14q32.11→qter of paternal origin encompassing PEGs of *DLK1* and *RTL1*, MEGs of *MEG3*, *RTL1as* and *MEG8*, and the differentially methylated regions (DMRs) of intergenic DMR (IG-DMR) and *MEG3*-DMR [37–40]. IG-DMR and *MEG3*-DMR, which are located between *DLK1* and *MEG3*, are respectively methylated on the paternal chromosome 14 and unmethylated on the maternal chromosome 14, and after paternal transmission, both DMRs are hypermethylated and after maternal transmission, both DMRs are hypomethylated [38,40–43]. Paternal uniparental disomy 14 [upd(14)pat] or Kagami–Ogata syndrome (OMIM 608149) is an imprinting disorder involving 14q32 imprinted genes and is characterized by skeletal abnormalities, joint contractures, dysmorphic facial features, developmental delay, mental retardation, a narrow and bell-shaped thorax with caudal bowing of the anterior ribs, cranial bowing of the posterior ribs, flaring of the iliac wings, abdominal wall defects, placentomegaly and polyhydramnios [44,45]. Maternal uniparental disomy 14 [upd(14)mat] or Temple syndrome (OMIM 616222) is an imprinting disorder involving 14q32 imprinted genes and is characterized by low birth weight, hypotonia, motor delay, feeding problems early in life, early puberty, a broad forehead, short nose, a wide nasal tip, small hands and feet [46,47]. Kagami–Ogata syndrome is caused by full or segmental upd(14)pat of normal chromosome, abnormal chromosome of Robertsonian 13q;14q translocation or i(14q), epimutations and microdeletions of 14q of maternal origin. The present case is not associated with




Figure 3. A karyotype of 46,XX,der(9)t(9;14)(p24.2;q32.11) in the fetus. The arrows indicate the breakpoints. der = derivative.




Figure 4. QF-PCR assays on the DNAs extracted from the placental tissue and the parental bloods using informative markers of D9S937 (9p24.2), D9S2149 (9p21.1), D14S140 (14q31.3), and D14S605 (14q32.2) show only one maternal allele at D9S937 (9p24.2) and a paternal allele:maternal allele ratio of 2:1 at D14S605 (14q32.2) in the fetus, indicating a paternal origin of the der(9) chromosome. QF-PCR = quantitative fluorescent polymerase chain reaction.

Kagami–Ogata syndrome. To date, no Kagami–Ogata syndrome has been reported in the case with partial trisomy 14q of paternal origin. Kagami–Ogata syndrome is associated with excessive *RTL1* expression with absent *MEG* expression or a ~2.5 times increased *RTL1* expression level with absence of functional *RTL1as*-encoded microRNAs [45].

In summary, we present molecular cytogenetic characterization of a prenatally detected der(9) of unknown origin which was confirmed to be der(9)t(9;14)(p24.2;q32.11). We discuss 9p deletion syndrome and 14q duplication syndrome in this case. Our presentation demonstrates the usefulness of molecular cytogenetic techniques such as aCGH, FISH, and QF-PCR in the determination of the origin and nature of a prenatally detected *de novo* derivative chromosome of unknown origin.

Conflict of interest

The authors have no conflicts of interest relevant to this article.

Acknowledgments

This work was supported by research grants MOST-103-2314-B-195-010 and MOST-104-2314-B-195-009 from the Ministry of Science and Technology and MMH-E-105-04 from Mackay Memorial Hospital, Taipei, Taiwan.

References

- [1] Chen CP, Lin MH, Chen YY, Chern SR, Chen YN, Wu PS, et al. Prenatal diagnosis and array comparative genomic hybridization characterization of interstitial deletions of 8q23.3–q24.11 and 8q24.13 associated with Langer-Giedion syndrome and Cornelia de Lange syndrome and haploinsufficiency of *TRPS1*, *RAD21* and *EXT1*. Taiwan J Obstet Gynecol 2015;54:592–6.
- [2] Chen CP, Su YN, Chen CY, Chern SR, Wu PS, Su JW, et al. Prenatal diagnosis and molecular cytogenetic characterization of a *de novo* pure distal 9p deletion and literature review. Genomics 2013;102:265–9.
- [3] Herpin A, Scharlt M. Sex determination: switch and suppress. Curr Biol 2011;21:R656–9.
- [4] Matson CK, Murphy MW, Sarver AL, Griswold MD, Bardwell VJ, Zarkower D. *DMRT1* prevents female reprogramming in the postnatal mammalian testis. Nature 2011;476:101–4.
- [5] Jørgensen A, Nielsen JE, Jensen MB, Græm N, Rajpert-De Meyts E. Analysis of meiosis regulators in human gonads: a sexually dimorphic spatio-temporal

- expression pattern suggests involvement of *DMRT1* in meiotic entry. Mol Hum Reprod 2012;18:523–34.
- [6] Ledig S, Hiort O, Wünsch L, Wieacker P. Partial deletion of *DMRT1* causes 46,XY ovotesticular disorder of sexual development. Eur J Endocrinol 2012;167:119–24.
- [7] Raymond CS, Shamu CE, Shen MM, Seifert KJ, Hirsch B, Hodgkin J, et al. Evidence for evolutionary conservation of sex-determining genes. Nature 1998;391:691–5.
- [8] Ying M, Chen B, Tian Y, Hou Y, Li Q, Shang X, et al. Nuclear import of human sexual regulator *DMRT1* is mediated by importin- β . Biochim Biophys Acta 2007;1773:804–13.
- [9] Griggs BL, Ladd S, Saul RA, DuPont BR, Srivastava AK. Dedicator of cytokinesis 8 is disrupted in two patients with mental retardation and developmental disabilities. Genomics 2008;91:195–202.
- [10] Lerer I, Sagi M, Meiner V, Cohen T, Zlotogora J, Abeliovich D. Deletion of the *ANKRD15* gene at 9p24.3 causes parent-of-origin-dependent inheritance of familial cerebral palsy. Hum Mol Genet 2005;14:3911–20.
- [11] Boycott KM, Flavelle S, Bureau A, Glass HC, Fujiwara TM, Wirrell E, et al. Homozygous deletion of the very low density lipoprotein receptor gene causes autosomal recessive cerebellar hypoplasia with cerebral gyral simplification. Am J Hum Genet 2005;77:477–83.
- [12] Moheb LA, Tzschach A, Garshabi M, Kahrizi K, Darvish H, Heshmati Y, et al. Identification of a nonsense mutation in the very low-density lipoprotein receptor gene (*VLDLR*) in an Iranian family with dysequilibrium syndrome. Eur J Hum Genet 2008;16:270–3.
- [13] Ozcelik T, Akarsu N, Uz E, Caglayan S, Gulsuner S, Onat OE, et al. Mutations in the very low-density lipoprotein receptor *VLDLR* cause cerebellar hypoplasia and quadrupedal locomotion in humans. Proc Natl Acad Sci USA 2008;105:4232–6.
- [14] Turkmen S, Hoffmann K, Demirhan O, Aruoba D, Humphrey N, Mundlos S. Cerebellar hypoplasia, with quadrupedal locomotion, caused by mutations in the very low-density lipoprotein receptor gene. Eur J Hum Genet 2008;16:1070–4.
- [15] Stumm M, Theile U, Kessel-Weiner E, Wieacker P, Pascu F, Ottolenghi C. Deletion of the DM-domain gene clusters in a fetus with ring chromosome 9 and sex reversal. Pediatr Pathol Mol Med 2000;19:415–23.
- [16] Vialard F, Ottolenghi C, Gonzales M, Choiset A, Girard S, Siffroi JP, et al. Deletion of 9p associated with gonadal dysfunction in 46,XY but not in 46,XX human fetuses. J Med Genet 2002;39:514–8.
- [17] Witters I, Vermeesch JR, Moerman PH, Fryns JP. Partial trisomy 3p/monosomy 9p with sex reversal. Ultrasound Obstet Gynecol 2004;23:418–9.
- [18] Brisset S, Kasakyan S, L'Herminé AC, Mairovitz V, Gautier E, Aubry M-C, et al. *De novo* monosomy 9p24.3-pter and trisomy 17p24.3-qter characterised by microarray comparative genomic hybridisation in a fetus with an increased nuchal translucency. Prenat Diagn 2006;26:206–13.
- [19] Chen CP, Lin CL, Lee CC, Wang W. Prenatal diagnosis of mosaic ring chromosome 9. Prenat Diagn 2006;26:870–1.
- [20] Chen CP, Su YN, Chern SR, Hsu CY, Tsai FJ, Wu PC, et al. Inv dup del(9p): prenatal diagnosis and molecular cytogenetic characterization by fluorescence *in situ* hybridization and array comparative genomic hybridization. Taiwan J Obstet Gynecol 2011;50:67–73.
- [21] Hou WC, Chen CP, Hwang KS, Chen YC, Lai YJ, Tien CY, et al. Prenatal diagnosis of a *de novo* 9p terminal chromosome deletion in a fetus with major congenital anomalies. Taiwan J Obstet Gynecol 2014;53:602–5.
- [22] Penacho V, Galán F, Martín-Bayón TA, Mayo S, Manchón I, et al. Prenatal diagnosis of a female fetus with ring chromosome 9, 46,XX,r(9)(p24q34), and a *de novo* interstitial 9p deletion. Cytogenet Genome Res 2014;144:275–9.
- [23] Orye E, Van Bever H, Desimpel H. Distal trisomy 14q due to tandem duplication (q24 leads to q32). Ann Génét 1983;26:238–9.
- [24] Turleau C, de Grouchy J, Chavin-Colin F, Denavit MF, Le Touze P. Distal 14q trisomy. Ann Génét 1983;26:165–70 [in French].
- [25] Mikelsaar RV, Ilus TA, Lurie IW. Distal trisomy 14q. J Med Genet 1987;24:380–1.
- [26] Chen CP, Chern SR, Lin SP, Lin CC, Li YC, Wang TH, et al. A paternally derived inverted duplication of distal 14q with a terminal 14q deletion. Am J Med Genet 2005;139A:146–50.
- [27] Thiel CT, Dörr HG, Trautmann U, Hoyer J, Hofmann K, Kraus C, et al. A *de novo* 7.6Mb tandem duplication of 14q32.2-qter associated with primordial short stature with neurosecretory growth hormone dysfunction, distinct facial anomalies and mild developmental delay. Eur J Med Genet 2008;51:362–7.
- [28] Trunca C, Optitz JM. Pericentric inversion of chromosome 14 and the risk of partial duplication of 14q (14q31–14qter). Am J Med Genet 1977;1:217–28.
- [29] Weinstein MR, Handmaker SD, Mohandas T. A unique pattern of dysmorphism and mental retardation associated with extra chromosome material (14q+): a probable duplication. Am J Hum Genet 1977;29:112A.
- [30] Sklower SL, Jenkins EC, Nolin SL, Duncan CJ, Warburton D, Anyane Yeboa K, et al. Distal duplication 14q: report of three cases and further delineation of the syndrome. Hum Genet 1984;68:159–64.
- [31] Carr DM, Jones-Quartey K, Vartanian MV, Moore-Kaplan H. Duplication 14 (q31–qter). J Med Genet 1987;24:372–4.
- [32] Wakita Y, Narahara K, Kikkawa K, Namba H, Hiramoto K, Eguchi K, et al. Distal 14q trisomy syndrome in two siblings: further delineation of its phenotype. Jpn J Hum Genet 1988;33:469–75.
- [33] Masada CT, Haskins Olney A, Fordyce R, Sanger WG. Partial deletion of 14q and partial duplication of 14q in sibs: testicular mosaicism for t(14q;14q) as a common mechanism. Am J Med Genet 1989;34:528–34.

- [34] Schinzel A. dup(14)(q31→qter). In: Schinzel A, editor. Catalogue of unbalanced chromosome aberrations in man. 2nd ed. Berlin: Water de Gruyter, GmbH & Co; 2001. p. 613–5.
- [35] Sonoda T, Kouno K, Sawada K, Sugimoto T. Distal 14q trisomy due to a maternal derivative chromosome 14. *Pediatr Int* 2001;43:313–5.
- [36] Chen CP, Hwang KS, Su HY, Lin SP, Su YN, Chern SR, et al. Prenatal diagnosis and molecular cytogenetic characterization of a *de novo* interstitial duplication of 14q (14q31.3→q32.12) associated with abnormal maternal serum biochemistry. *Taiwan J Obstet Gynecol* 2013;52:125–8.
- [37] Hosoki K, Ogata T, Kagami M, Tanaka T, Saitoh S. Epimutation (hypomethylation) affecting the chromosome 14q32.2 imprinted region in a girl with upd(14)mat-like phenotype. *Eur J Hum Genet* 2008;16:1019–23.
- [38] Kagami M, Sekita Y, Nishimura G, Irie M, Kato F, Okada M, et al. Deletions and epimutations affecting the human 14q32.2 imprinted region in individuals with paternal and maternal upd(14)-like phenotypes. *Nat Genet* 2008;40:237–42.
- [39] Kagami M, O'Sullivan MJ, Green AJ, Watabe Y, Arisaka O, Masawa N, et al. The IG-DMR and the MEG3-DMR at human chromosome 14q32.2: hierarchical interaction and distinct functional properties as imprinting control centers. *PLoS Genet* 2010;6:e1000992.
- [40] Ogata T, Kagami M, Ferguson-Smith AC. Molecular mechanisms regulating phenotypic outcome in paternal and maternal uniparental disomy for chromosome 14. *Epigenetics* 2008;3:181–7.
- [41] Paulsen M, Takada S, Youngson NA, Benchaib M, Charlier C, Segers K, et al. Comparative sequence analysis of the imprinted Dlk1-Gtl2 locus in three mammalian species reveals highly conserved genomic elements and refines comparison with the Igf2-H19 region. *Genome Res* 2001;11:2085–94.
- [42] Lin SP, Youngson N, Takada S, Seitz H, Reik W, Paulsen M, et al. Asymmetric regulation of imprinting on the maternal and paternal chromosomes at the Dlk1-Gtl2 imprinted cluster on mouse chromosome 12. *Nat Genet* 2003;35:97–102.
- [43] Beygo J, Elbracht M, de Groot K, Begemann M, Kanber D, Platzer K, et al. Novel deletions affecting the MEG3-DMR provide further evidence for a hierarchical regulation of imprinting in 14q32. *Eur J Hum Genet* 2015;23:180–8.
- [44] Kagami M, Kurosawa K, Miyazaki O, Ishino F, Matsuoka K, Ogata T. Comprehensive clinical studies in 34 patients with molecularly defined UPD(14)pat and related conditions (Kagami-Ogata syndrome). *Eur J Hum Genet* 2015;23:1488–98.
- [45] Ogata T, Kagami M. Kagami-Ogata syndrome: a clinically recognizable upd(14)pat and related disorder affecting the chromosome 14q32.2 imprinted region. *J Hum Genet* 2016;61:87–94.
- [46] Temple IK, Cockwell A, Hassold T, Pettay D, Jacobs P. Maternal uniparental disomy for chromosome 14. *J Med Genet* 1991;28:511–4.
- [47] Temple IK, Shrubbs V, Lever M, Bullman H, Mackay DJG. Isolated imprinting mutation of the *DLK1/GTL2* locus associated with a clinical presentation of maternal uniparental disomy of chromosome 14. *J Med Genet* 2007;44:637–40.